



# Many-Flavor Domain Wall Fermions and Fixed Topology

Meifeng Lin



New Horizons for Lattice Computations with Chiral Fermions  
BNL, May 14 - 16, 2012



# Outline

## Motivation

## LSD Calculations with Domain Wall Fermions

Recap: Results for  $N_f = 2, 6$

New: Results for  $N_f = 10$

## Summary



## Motivation

- ▶ SU(N) gauge theories with many flavors of fermions may exhibit behaviors that are fundamentally different from QCD.
- ▶ Increasing  $N_f$  causes the gauge coupling to run slowly, and “walking” may occur, which could lead to some properties desirable for Technicolor model building.
- ▶ Specifically, we are interested in
  - ▶ enhanced chiral condensate
  - ▶ small  $S$  parameter
  - ▶ new pattern in hadron spectrum
- ▶ When  $N_f > N_f^c$ , the theory becomes conformal, and spectrum vanishes at the chiral limit. Also on our wishlist is to
  - ▶ locate  $N_f^c$  non-perturbatively



# The Lattice Strong Dynamics Collaboration

Tom Appelquist Yale U.  
Ron Babich NVIDIA  
Rich Brower Boston U.  
Michael Buchoff LLNL  
Michael Cheng Boston U.  
Mike Clark NVIDIA  
Saul Cohen U. Washington  
George Fleming Yale U.  
Joe Kiskis U. of California, Davis  
Meifeng Lin Yale U.

Heechang Na ANL  
Ethan Neil Fermilab  
James Osborn ANL  
Claudio Rebbi Boston U.  
David Schaich Boston U.  
Chris Schroeder LLNL  
Sergey Syritsyn LBL  
Joe Wasem LLNL  
Oliver Witzel Boston U.  
Gennady Voronov Yale U.  
Pavlos Vranas LLNL



## Gauge Ensembles

- ▶ **Fermion action:** domain wall fermions.
- ▶ **Gauge action:** Iwasaki.
- ▶ **Ensemble size:** Typically  $\sim 1200$  MD trajectories.

[Cost grows as  $N_f^{3/2}$ ]

$N_f$	Volume	$L_s$	$\beta$	$am_f$	$am_{\text{res}}$
2	$32^3 \times 64$	16	2.70	0.005...0.030	$2.5 \times 10^{-5}$
6	$32^3 \times 64$	16	2.10	0.005...0.030	$8.2 \times 10^{-4}$
10	$32^3 \times 64$	16	1.95	0.005...0.030	$1.7 \times 10^{-3}$


- ▶ For  $N_f = 2$ ,  $aM_V \approx 0.21 \Rightarrow a^{-1} \approx 3.7$  GeV.
- ▶ For  $N_f = 6$  and 10, match  $aM_V$  to within 10% of  $N_f = 2$ .



Recap: Results for  $N_f = 2, 6$

# Condensate Enhancement


$$R_m \equiv \frac{M_m^2}{2mF_\pi(N_f=6)} / \frac{M_m^2}{2mF_\pi(N_f=2)}$$



T. Appelquist *et al.* (LSD Collaboration), PRL104:071601,2010

Recap: Results for  $N_f = 2, 6$

# The S Parameter





T. Appelquist *et al.* (LSD Collaboration), PRL 106 (2011) 231601



New: Results for  $N_f = 10$

## A New Observation for $N_f = 10$

- ▶ Two independent ensembles for each quark mass, with **ordered** and **disordered** start.
- ▶ The ensembles thermalize to different values
- ▶ **NOT** observed for  $N_f = 2$  or 6.






○ ○  
 ○ ● ○ ○ ○ ○ ○ ○

New: Results for  $N_f = 10$

## Starting-Config. Dependence

Statistically significant difference between the ordered-start and disordered-start ensembles are observed for almost all the ensembles.



New: Results for  $N_f = 10$ 

## Slowly-moving/Fixed Topology

$m_f$	start	trajectory range	$Q$
0.005	O	500 - 1319	0
0.005	D	500 - 919	3
0.010	O	500 - 1373	0
0.010	D	500 - 1259	1
0.015	O	500 - 1204	0
0.015	D	500 - 1450	-1
0.020	O	500 - 1239	0
0.020	D	500 - 1221	12
0.025	O	500 - 1256	0
0.025	D	500 - 609	-19
		610 - 942	-18
		943 - 1095	-17
		1096-1097	-16
		1098 - 1382	-17
		1383 - 1484	-16
0.030	O	500 - 1415	0
0.030	D	500 - 1227	9



New: Results for  $N_f = 10$

## Effects of Fixed Topology

- ▶ The slight topological tunneling at  $m_f = 0.025$  (D) allows us to analyze the data for different topological sectors separately.
- ▶ Given the limited number of measurements for each topology, we only block every trajectory, hence the errors will be underestimated.
- ▶ Nevertheless, such analysis will give us a crude idea how much the fixed topology may affect our results.

R. Brower, S. Chandrasekharan, John W. Negele, and U.J. Wiese. 2003 has the following formula for the  $Q$ -dependence


$$M_Q = M(0) + \frac{1}{2} M^{(2)}(0) \frac{1}{V_{\chi_t}} \left( 1 - \frac{Q^2}{V_{\chi_t}} \right) + O\left(\frac{1}{V^3}\right).$$

$\chi_t$  is the topological susceptibility,  $M^{(2)}(0)$  is the second derivative with respect to  $\theta$  at  $\theta = 0$ , and  $M(0)$  is the mass at  $\theta = 0$  (or equivalently, at non-fixed  $Q$ ).



New: Results for  $N_f = 10$

## Observed $Q$ -Dependence



- ▶ The discrepancy between ordered and disordered ensembles can be described by the dependence on the fixed topology.
- ▶ To get results at  $\theta = 0$ , we still need to know the **topological susceptibility**  $\rightarrow$  **work in progress**.
- ▶ Large discrepancies  $\Rightarrow$  small  $\chi_t$ ?  $V$  too small?



New: Results for  $N_f = 10$

## Combining Datasets

Lacking the results for the topological susceptibility, can we still combine the two ensembles and obtain some best estimates of the "true" value,  $M(0)$ ?

- ▶ From our fits,  $M^{(2)}(0)$  is negative, so  $M(0)$  is between the results for  $Q = 0$  and  $Q \neq 0$ .
- ▶ Take  $\overline{M} = (M_0 + M_Q)/2$  as the central value.
- ▶ Take the largest relative difference between  $M_Q$  and  $M_0$  as systematic error.
- ▶ Combine statistical and systematic errors as final error.




New: Results for  $N_f = 10$

## Testing Conformality - Qualitatively

In a conformal theory with a small finite mass,

$$M_X = C_X m^{1/(1+\gamma^*)}$$



so the vector-to-pseudoscalar meson mass ratio  $M_V/M_P$  should look roughly constant.



- ▶ The  $N_f = 2$  data are diverging as  $m \rightarrow 0$ , as expected since  $M_P \rightarrow 0$ .
- ▶ Similarly for  $N_f = 6$ , though not as fast as  $N_f = 2$ .
- ▶ For  $N_f = 10$  the ratios seem to be roughly constant.

New: Results for  $N_f = 10$ 

# Testing Conformality - Quantitatively



Obs.	$m_f \geq 0.010$	$m_f \geq 0.015$	$m_f \geq 0.020$
$\gamma^*$	1.45(15)	0.94(16)	1.18(44)
[68% CI]	[0.88, 2.47]	[0.68, 1.44]	[1.01, 1.69]
$C_P$	0.978(9)	1.44(21)	1.21(42)
$C_V$	1.168(10)	1.70(25)	1.42(49)
$C_A$	1.429(13)	2.14(32)	1.79(63)
$C_N$	1.749(16)	2.53(37)	2.10(73)
$C_{N^*}$	2.232(25)	3.35(56)	2.87(1.02)
$C_{FP}$	0.121(12)	0.190(29)	0.164(57)
$\chi^2/\text{d.o.f.}$	69/31	14/23	3/15

Finite volume effects may be large for  
 $am_f \leq 0.01$ .



## Summary

- ▶ First SU(3) lattice simulations with  $N_f = 6$  and 10 domain wall fermions have been performed.
- ▶ Hints of condensate enhancement and reduced  $S$  parameter have been seen with  $N_f = 6$ .
- ▶ With increasing  $N_f$ , the topology becomes more difficult to evolve. Effects of fixed topology are big for  $N_f = 10$ .
- ▶ Spectrum for  $N_f = 10$  is consistent with conformality. **But cannot rule out the chirally-broken scenario, as the masses are too heavy to make use of ChPT.**

## Future Work

- ▶ Calculate topological susceptibility at fixed  $Q$  for  $N_f = 10$ .  
*S. Aoki et al., 2007, 2008*
- ▶ Quantative studies of finite volume effects
- ▶ Smaller quark masses